

TABLE 1.

TURFGRASS COLOR RATINGS FOR 2004 IN ITHACA, NY
FOLLOWING THE APPLICATION OF WETTING AGENTS.

TURFGRASS COLOR RATINGS ^a										
NAME	1 WEEK ^b	3 WEEKS	5 WEEKS	7 WEEKS	9 WEEKS	11 WEEKS	13 WEEKS	15 WEEKS	17 WEEKS	YEARLY AVE.
AQUEDUCT	7.0	7.0	7.0	7.0	8.0	7.0	7.0	7.0	8.0	7.2
BRILLIANCE	7.5	7.3	7.0	8.0	8.0	8.0	7.5	8.0	9.0	7.8
CASCADE PLUS	7.0	6.5	6.5	7.0	7.5	7.0	7.0	7.0	8.0	7.1
CONTROL	7.5	7.3	7.0	7.5	7.5	7.5	7.5	7.5	8.5	7.5
HYDRO-WET	7.0	7.0	7.0	7.5	8.0	7.5	7.0	7.5	8.5	7.4
LESCOFLO	7.5	7.3	7.0	7.5	7.5	7.5	7.5	7.5	8.5	7.5
NAIAD	7.0	7.0	7.0	7.0	8.0	7.0	7.0	7.0	8.0	7.2
PRIMER SELECT	7.3	7.3	7.3	8.0	8.3	7.8	7.3	8.0	9.0	7.8
RESPOND 2	7.0	7.0	6.8	7.5	7.8	7.5	7.0	7.5	8.5	7.4
SURFSIDE 37	7.0	6.8	6.5	7.0	7.0	7.0	7.0	7.0	8.0	7.0
TRICURE	7.3	7.0	7.0	7.5	7.5	7.3	7.3	7.5	8.5	7.4
LSD ^c	0.8	0.8	0.6	0.6	0.7	0.7	0.8	0.6	0.6	0.5
CV ^d (%)	5.3	5.5	4.8	5.3	5.3	5.6	5.3	5.3	4.6	4.1

a) Turfgrass color was visually rated using a scale of 1=brown, 5=medium green and 9=dark green.

b) Color ratings were taken every two weeks beginning one week after the initial wetting agent application which was made on June 5, 2004.

c) LSD is the least significant difference among the treatment means. To determine if one treatment is significantly different from another, subtract the mean of one treatment from the mean of another treatment. A statistically significant difference occurs when this value is larger than the LSD value given at the bottom of the column. Treatment means should be compared only within a column (LSD 0.05).

d) CV is the coefficient of variation and indicates the percent variation of the treatment means in each column.

TABLE 2.

TURFGRASS QUALITY RATINGS FOR 2004 IN ITHACA, NY
FOLLOWING THE APPLICATION OF WETTING AGENTS.

TURFGRASS QUALITY RATINGS ^a										YEARLY AVE.
NAME	1 WEEK ^b	3 WEEKS	5 WEEKS	7 WEEKS	9 WEEKS	11 WEEKS	13 WEEKS	15 WEEKS	17 WEEKS	
AQUEDUCT	8.0	7.0	7.0	7.0	7.0	7.0	7.0	7.0	7.0	7.1
BRILLIANCE	8.3	8.0	7.5	7.0	7.5	7.8	7.3	8.0	7.5	7.6
CASCADE PLUS	7.5	7.0	7.0	6.5	7.0	7.0	6.5	7.0	7.0	6.9
CONTROL	8.3	7.5	7.5	6.5	7.5	7.5	7.3	7.5	7.5	7.4
HYDRO-WET	8.0	7.5	7.0	7.0	7.0	7.0	7.0	7.5	7.0	7.2
LESCOFLO	8.3	7.5	7.5	6.5	7.5	7.5	7.3	7.5	7.5	7.4
NAIAD	8.0	7.0	7.0	7.0	7.0	7.0	7.0	7.0	7.0	7.1
PRIMER SELECT	8.3	8.0	7.3	7.0	7.3	7.8	7.3	8.0	7.3	7.6
RESPOND 2	7.8	7.5	7.0	6.8	7.0	7.3	7.0	7.5	7.0	7.2
SURFSIDE 37	7.8	7.0	7.0	6.0	7.0	7.0	6.8	7.0	7.0	6.9
TRICURE	8.0	7.5	7.3	6.5	7.3	7.3	7.0	7.5	7.3	7.3
LSD ^c	0.8	0.6	0.8	0.6	0.8	0.7	0.8	0.6	0.8	0.5
CV ^d (%)	5.1	5.3	5.3	5.4	5.3	5.5	5.5	5.3	5.3	4.2

- a) Turfgrass quality was visually rated using a scale of 1=poor quality, 5=acceptable quality and 9=excellent quality.
- b) Turfgrass quality ratings were taken every two weeks beginning one week after the initial wetting agent application which was made on June 5, 2004.
- c) LSD is the least significant difference among the treatment means. To determine if one treatment is significantly different from another, subtract the mean of one treatment from the mean of another treatment. A statistically significant difference occurs when this value is larger than the LSD value given at the bottom of the column. Treatment means should be compared only within a column (LSD 0.05).
- d) CV is the coefficient of variation and indicates the percent variation of the treatment means in each column.

TABLE 3.

PHYTOTOXICITY RATINGS FOR 2004 IN ITHACA, NY
 FOLLOWING THE APPLICATION OF WETTING AGENTS.
 THE INITIAL APPLICATION OF WETTING AGENTS WAS MADE ON JUNE 5, 2004.

PHYTOTOXICITY RATINGS^a

NAME	APP 1-1 ^b	APP 1-3	APP 1-7	APP 2-1	APP 2-3	APP 2-7	APP 3-1	APP 3-3	APP 3-7	APP 4-1	APP 4-3	APP 4-7	APP 5-1	APP 5-3	APP 5-7
AQUEDUCT	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
BRILLIANCE	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
CASCADE PLUS	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
CONTROL	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
HYDRO-WET	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
LESCOFLO	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
NAIAD	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
PRIMER SELECT	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
RESPOND 2	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
SURFSIDE 37	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
TRICURE	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
LSD ^c	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
CV ^d (%)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

a) Phytotoxicity was visually rated using a scale of 1=brown or discolored turf, 7=acceptable damage and 9=green turf, no damage.

b) Phytotoxicity ratings were taken one, three and seven days after each application of any wetting agent. App 1-1 refers to application number one, the ratings were taken one day after application.

c) LSD is the least significant difference among the treatment means. To determine if one treatment is significantly different from another, subtract the mean of one treatment from the mean of another treatment. A statistically significant difference occurs when this value is larger than the LSD value given at the bottom of the column. Treatment means should be compared only within a column (LSD 0.05).

d) CV is the coefficient of variation and indicates the percent variation of the treatment means in each column.

TABLE 3. (CONTINUED):

PHYTOTOXICITY RATINGS FOR 2004 IN ITHACA, NY
 FOLLOWING THE APPLICATION OF WETTING AGENTS.
 THE INITIAL APPLICATION OF WETTING AGENTS WAS MADE ON JUNE 5, 2004.

PHYTOTOXICITY RATINGS ^a															
NAME	APP 6-1 ^b	APP 6-3	APP 6-7	APP 7-1	APP 7-3	APP 7-7	APP 8-1	APP 8-3	APP 8-7	APP 9-1	APP 9-3	APP 9-7	APP 10-1	APP 10-3	APP 10-7
AQUEDUCT	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
BRILLIANCE	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
CASCADE PLUS	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
CONTROL	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
HYDRO-WET	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
LESCOFLO	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
NAIAD	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
PRIMER SELECT	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
RESPOND 2	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
SURFSIDE 37	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
TRICURE	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
LSD ^c	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
CV ^d (%)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

- a) Phytotoxicity was visually rated using a scale of 1=brown or discolored turf, 7=acceptable damage and 9=green turf, no damage.
- b) Phytotoxicity ratings were taken one, three and seven days after each application of any wetting agent. App 6-1 refers to application number six, the ratings were taken one day after application.
- c) LSD is the least significant difference among the treatment means. To determine if one treatment is significantly different from another, subtract the mean of one treatment from the mean of another treatment. A statistically significant difference occurs when this value is larger than the LSD value given at the bottom of the column. Treatment means should be compared only within a column (LSD 0.05).
- d) CV is the coefficient of variation and indicates the percent variation of the treatment means in each column.

TABLE 3. (CONTINUED):

PHYTOTOXICITY RATINGS FOR 2004 IN ITHACA, NY
 FOLLOWING THE APPLICATION OF WETTING AGENTS.
 THE INITIAL APPLICATION OF WETTING AGENTS WAS MADE ON JUNE 5, 2004.

PHYTOTOXICITY RATINGS^a

NAME	APP 11-1 ^b	APP 11-3	APP 11-7	APP 12-1	APP 12-3	APP 12-7	APP 13-1	APP 13-3	APP 13-7	APP 14-1	APP 14-3	APP 14-7	APP 15-1	APP 15-3	APP 15-7
AQUEDUCT	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
BRILLIANCE	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
CASCADE PLUS	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
CONTROL	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
HYDRO-WET	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
LESCOFLO	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
NAIAD	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
PRIMER SELECT	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
RESPOND 2	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
SURFSIDE 37	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
TRICURE	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
LSD ^c	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
CV ^d (%)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

a) Phytotoxicity was visually rated using a scale of 1=brown or discolored turf, 7=acceptable damage and 9=green turf, no damage.

b) Phytotoxicity ratings were taken one, three and seven days after each application of any wetting agent. App 11-1 refers to application number 11, the ratings were taken one day after application.

c) LSD is the least significant difference among the treatment means. To determine if one treatment is significantly different from another, subtract the mean of one treatment from the mean of another treatment. A statistically significant difference occurs when this value is larger than the LSD value given at the bottom of the column. Treatment means should be compared only within a column (LSD 0.05).

d) CV is the coefficient of variation and indicates the percent variation of the treatment means in each column.

TABLE 4. WATER DROPLET PENETRATION TIMES BY DEPTH FOR 2004 IN ITHACA, NY
TWO WEEKS AFTER THE INITIAL APPLICATION OF WETTING AGENTS.
THE INITIAL APPLICATION OF WETTING AGENTS WAS MADE ON JUNE 5, 2004.

WATER DROPLET PENETRATION MEASURED IN SECONDS ^a						
NAME	0.5 CM ^b	1.5 CM	2.5 CM	3.5 CM	4.5 CM	5.5 CM
AQUEDUCT	1.5	7.8	5.0	0.5	0.0	0.0
BRILLIANCE	1.0	1.3	0.8	0.8	0.0	0.0
CASCADE PLUS	1.0	1.5	1.5	0.3	0.0	0.0
CONTROL	60.8	49.8	3.3	0.8	0.0	0.0
HYDRO-WET	29.3	35.3	6.5	2.3	0.8	0.0
LESCOFLO	14.3	20.3	9.3	1.8	0.3	0.0
NAIAD	146.5	57.5	9.5	2.8	0.5	0.0
PRIMER SELECT	9.3	11.0	6.0	1.0	0.3	0.0
RESPOND 2	86.5	32.5	8.5	4.5	5.3	0.8
SURFSIDE 37	67.0	49.5	15.0	1.8	0.3	0.3
TRICURE	3.0	12.0	3.8	1.5	0.8	0.0
LSD ^c	26.3	42.1	8.0	5.5	7.5	1.1
CV ^d (%)	52.4	98.8	77.2	148.5	438.7	529.6

a) The maximum time for water droplet penetration was 600 seconds. Any water droplet remaining after 600 seconds was recorded as 600 seconds.

b) Depth in centimeters below the soil surface.

c) LSD is the least significant difference among the treatment means. To determine if one treatment is significantly different from another, subtract the mean of one treatment from the mean of another treatment. A statistically significant difference occurs when this value is larger than the LSD value given at the bottom of the column. Treatment means should be compared only within a column (LSD 0.05).

d) CV is the coefficient of variation and indicates the percent variation of the treatment means in each column.

TABLE 5. WATER DROPLET PENETRATION TIMES BY DEPTH FOR 2004 IN ITHACA, NY
 FOUR WEEKS AFTER THE INITIAL APPLICATION OF WETTING AGENTS.
 THE INITIAL APPLICATION OF WETTING AGENTS WAS MADE ON JUNE 5, 2004.

WATER DROPLET PENETRATION MEASURED IN SECONDS ^a						
NAME	0.5 CM ^b	1.5 CM	2.5 CM	3.5 CM	4.5 CM	5.5 CM
AQUEDUCT	7.8	5.3	0.3	0.0	0.0	0
BRILLIANCE	4.8	2.3	0.5	0.5	0.0	0
CASCADE PLUS	9.8	4.5	0.8	0.3	0.3	0
CONTROL	47.5	15.8	0.8	0.0	0.0	0
HYDRO-WET	23.5	11.0	2.8	0.0	0.0	0
LESCOFLO	17.0	6.8	0.3	0.0	0.0	0
NAIAD	57.0	16.8	2.8	0.0	0.0	0
PRIMER SELECT	18.0	8.5	0.8	0.0	0.0	0
RESPOND 2	49.5	11.0	0.8	0.0	0.0	0
SURFSIDE 37	37.8	10.5	1.0	0.0	0.0	0
TRICURE	9.5	4.0	0.8	0.0	0.3	0
LSD ^c	13.0	12.7	3.9	0.4	0.5	0
CV ^d (%)	38.2	77.7	167.5	328.0	476.8	.

a) The maximum time for water droplet penetration was 600 seconds. Any water droplet remaining after 600 seconds was recorded as 600 seconds.

b) Depth in centimeters below the soil surface.

c) LSD is the least significant difference among the treatment means. To determine if one treatment is significantly different from another, subtract the mean of one treatment from the mean of another treatment. A statistically significant difference occurs when this value is larger than the LSD value given at the bottom of the column. Treatment means should be compared only within a column (LSD 0.05).

d) CV is the coefficient of variation and indicates the percent variation of the treatment means in each column.

TABLE 6. WATER DROPLET PENETRATION TIMES BY DEPTH FOR 2004 IN ITHACA, NY
EIGHT WEEKS AFTER THE INITIAL APPLICATION OF WETTING AGENTS.
THE INITIAL APPLICATION OF WETTING AGENTS WAS MADE ON JUNE 5, 2004.

WATER DROPLET PENETRATION MEASURED IN SECONDS ^a						
NAME	0.5 CM ^b	1.5 CM	2.5 CM	3.5 CM	4.5 CM	5.5 CM
AQUEDUCT	5.3	5.5	0.3	0.0	0.0	0.0
BRILLIANCE	12.0	7.5	2.0	0.0	0.0	0.0
CASCADE PLUS	20.5	7.5	8.5	2.8	1.3	0.8
CONTROL	6.5	7.5	5.3	5.8	0.5	0.0
HYDRO-WET	24.0	15.5	0.5	0.3	0.3	0.0
LESCOFLO	23.5	13.0	1.0	0.3	0.0	0.0
NAIAD	44.0	35.0	5.5	0.8	0.3	0.3
PRIMER SELECT	14.5	10.5	1.5	0.5	0.5	0.0
RESPOND 2	40.0	15.0	2.0	0.3	0.3	0.0
SURFSIDE 37	16.5	7.3	0.5	0.0	0.0	0.0
TRICURE	6.0	6.5	1.3	0.0	0.0	0.0
LSD ^c	36.3	15.5	14.1	5.4	2.2	1.1
CV ^d (%)	98.9	80.3	228.1	288.8	326.1	508.4

- a) The maximum time for water droplet penetration was 600 seconds. Any water droplet remaining after 600 seconds was recorded as 600 seconds.
- b) Depth in centimeters below the soil surface.
- c) LSD is the least significant difference among the treatment means. To determine if one treatment is significantly different from another, subtract the mean of one treatment from the mean of another treatment. A statistically significant difference occurs when this value is larger than the LSD value given at the bottom of the column. Treatment means should be compared only within a column (LSD 0.05).
- d) CV is the coefficient of variation and indicates the percent variation of the treatment means in each column.

TABLE 7. WATER DROPLET PENETRATION TIMES BY DEPTH FOR 2004 IN ITHACA, NY
 12 WEEKS AFTER THE INITIAL APPLICATION OF WETTING AGENTS.
 THE INITIAL APPLICATION OF WETTING AGENTS WAS MADE ON JUNE 5, 2004.

WATER DROPLET PENETRATION MEASURED IN SECONDS ^a						
NAME	0.5 CM ^b	1.5 CM	2.5 CM	3.5 CM	4.5 CM	5.5 CM
AQUEDUCT	7.0	5.8	1.0	0.0	0.0	0.0
BRILLIANCE	18.5	7.0	2.8	0.0	0.0	0.0
CASCADE PLUS	20.3	13.8	2.5	0.5	0.0	0.0
CONTROL	29.3	12.0	0.5	0.0	0.0	0.0
HYDRO-WET	17.8	21.8	2.5	0.0	0.0	0.0
LESCOFLO	34.5	10.0	1.0	0.0	0.0	0.0
NAIAD	50.0	25.3	4.8	0.3	0.0	0.0
PRIMER SELECT	17.0	9.0	0.5	0.0	0.0	0.0
RESPOND 2	35.3	10.0	1.0	0.0	0.0	0.0
SURFSIDE 37	37.0	13.3	2.8	0.0	0.0	0.0
TRICURE	27.3	9.5	2.0	1.0	0.3	0.5
LSD ^c	29.3	9.1	4.0	1.7	0.3	0.7
CV ^d (%)	61.1	48.1	105.1	438.4	663.3	663.3

a) The maximum time for water droplet penetration was 600 seconds. Any water droplet remaining after 600 seconds was recorded as 600 seconds.

b) Depth in centimeters below the soil surface.

c) LSD is the least significant difference among the treatment means. To determine if one treatment is significantly different from another, subtract the mean of one treatment from the mean of another treatment. A statistically significant difference occurs when this value is larger than the LSD value given at the bottom of the column. Treatment means should be compared only within a column (LSD 0.05).

d) CV is the coefficient of variation and indicates the percent variation of the treatment means in each column.

TABLE 8. WATER DROPLET PENETRATION TIMES BY DEPTH FOR 2004 IN ITHACA, NY
 16 WEEKS AFTER THE INITIAL APPLICATION OF WETTING AGENTS.
 THE INITIAL APPLICATION OF WETTING AGENTS WAS MADE ON JUNE 5, 2004.

WATER DROPLET PENETRATION MEASURED IN SECONDS ^a						
NAME	0.5 CM ^b	1.5 CM	2.5 CM	3.5 CM	4.5 CM	5.5 CM
AQUEDUCT	9.3	5.8	0.3	0.0	0.0	0
BRILLIANCE	21.5	16.5	1.0	0.0	0.0	0
CASCADE PLUS	28.5	11.0	0.3	0.0	0.0	0
CONTROL	39.3	22.5	9.8	3.8	0.8	0
HYDRO-WET	12.8	6.8	0.5	0.3	0.0	0
LESCOFLO	35.0	17.3	5.0	0.0	0.0	0
NAIAD	68.5	15.8	0.3	0.0	0.0	0
PRIMER SELECT	27.3	7.0	0.5	0.0	0.0	0
RESPOND 2	47.5	16.3	3.5	0.0	0.0	0
SURFSIDE 37	47.3	9.8	0.5	0.0	0.0	0
TRICURE	11.8	6.8	0.5	0.0	0.0	0
LSD ^c	25.3	17.6	7.2	1.6	1.0	0
CV ^d (%)	53.6	71.9	201.8	287.4	663.3	.

- a) The maximum time for water droplet penetration was 600 seconds. Any water droplet remaining after 600 seconds was recorded as 600 seconds.
- b) Depth in centimeters below the soil surface.
- c) LSD is the least significant difference among the treatment means. To determine if one treatment is significantly different from another, subtract the mean of one treatment from the mean of another treatment. A statistically significant difference occurs when this value is larger than the LSD value given at the bottom of the column. Treatment means should be compared only within a column (LSD 0.05).
- d) CV is the coefficient of variation and indicates the percent variation of the treatment means in each column.

TABLE 9. YEARLY AVERAGE WATER DROPLET PENETRATION TIMES BY DEPTH FOR 2004 IN ITHACA, NY AFTER THE APPLICATION OF WETTING AGENTS.

WATER DROPLET PENETRATION MEASURED IN SECONDS ^a						
NAME	0.5 CM ^b	1.5 CM	2.5 CM	3.5 CM	4.5 CM	5.5 CM
AQUEDUCT	6.2	6.0	1.4	0.1	0.0	0.0
BRILLIANCE	11.6	6.9	1.4	0.3	0.0	0.0
CASCADE PLUS	16.0	7.7	2.7	0.8	0.3	0.2
CONTROL	36.7	21.5	3.9	2.1	0.3	0.0
HYDRO-WET	21.5	18.1	2.6	0.6	0.2	0.0
LESCOFLO	24.9	13.5	3.3	0.4	0.1	0.0
NAIAD	73.2	30.1	4.6	0.8	0.2	0.1
PRIMER SELECT	17.2	9.2	1.9	0.3	0.2	0.0
RESPOND 2	51.8	17.0	3.2	1.0	1.1	0.2
SURFSIDE 37	41.1	18.1	4.0	0.4	0.1	0.1
TRICURE	11.5	7.8	1.7	0.5	0.3	0.1
LSD ^c	8.6	10.4	4.8	1.4	1.5	0.3
CV ^d (%)	23.2	49.7	77.3	120.5	287.0	287.6

- a) The maximum time for water droplet penetration was 600 seconds. Any water droplet remaining after 600 seconds was recorded as 600 seconds.
- b) Depth in centimeters below the soil surface.
- c) LSD is the least significant difference among the treatment means. To determine if one treatment is significantly different from another, subtract the mean of one treatment from the mean of another treatment. A statistically significant difference occurs when this value is larger than the LSD value given at the bottom of the column. Treatment means should be compared only within a column (LSD 0.05).
- d) CV is the coefficient of variation and indicates the percent variation of the treatment means in each column.

TABLE 10. DEW FORMATION/CONTROL AND PERCENT ALGAE RATINGS FOR 2004 IN ITHACA, NY FOLLOWING THE APPLICATION OF WETTING AGENTS.

DEW FORMATION/CONTROL AND PERCENT ALGAE RATINGS^a

NAME	DEW 1 ^b	DEW 2	DEW 3	YEARLY AVE.	PERCENT ALGAE ^b
AQUEDUCT	1	1	3.0	1.7	17.5
BRILLIANCE	1	1	1.0	1.0	12.0
CASCADE PLUS	1	1	1.3	1.1	21.3
CONTROL
HYDRO-WET	1	1	1.5	1.2	21.8
LESCOFLO	1	1	1.0	1.0	5.0
NAIAD	1	1	3.8	1.9	15.0
PRIMER SELECT	1	1	4.8	2.3	15.0
RESPOND 2	1	1	5.0	2.3	6.3
SURFSIDE 37	1	1	3.0	1.7	11.3
TRICURE	1	1	4.3	2.1	25.0
LSD ^c	0	0	2.1	0.7	24.3
CV ^d (%)	0	0	49.0	28.8	76.9

- a) Dew formation/control was visually rated using a scale of 1=heavy dew present and 9= no dew present. Algae was rated as percent of plot covered.
- b) Dew formation/control ratings and algae ratings were taken on various dates following the application of wetting agents.
- c) LSD is the least significant difference among the treatment means. To determine if one treatment is significantly different from another, subtract the mean of one treatment from the mean of another treatment. A statistically significant difference occurs when this value is larger than the LSD value given at the bottom of the column. Treatment means should be compared only within a column (LSD 0.05).
- d) CV is the coefficient of variation and indicates the percent variation of the treatment means in each column.